

## **REFERENCES AND SOURCES: SUBSTANCE MISUSE**

Abdulrahim, D. (2001). Substance Misuse and Mental Health Co-morbidity (Dual Diagnosis). London: Health Advisory Service.

[http://handbooks.homeless.org.uk/hostels/individuals/therapeutic/Dual\\_Diagnosis\\_NTA.pdf/view?searchterm=mental](http://handbooks.homeless.org.uk/hostels/individuals/therapeutic/Dual_Diagnosis_NTA.pdf/view?searchterm=mental)

Anastasiadou, D., Sepulveda, A.R., Sanchez, J.C., Parks, M., Alvarez, T. & Graell, M. (2016). Family functioning and quality of life among families in eating disorders: A comparison with substance-related disorders and healthy controls. European Eating Disorders Review, 24(4), 294-303.

Baker, S. (2000). Trouble Within – Alcohol, Children and Families. Conference Report. Alcohol Concern. London: Alcohol Concern.

[http://www.alcoholconcern.org.uk/files/20030904\\_142841\\_Trouble%20within%20June%202000.prn.pdf](http://www.alcoholconcern.org.uk/files/20030904_142841_Trouble%20within%20June%202000.prn.pdf)

Barrowclough, C., Haddock, G., Tarrier, N., Lewis, S.W., Moring, J., O'Brien, R., Schofield, N. & McGovern, J. (2001). Randomized Controlled Trial of Motivational Interviewing, Cognitive Behavior Therapy, and Family Intervention for Patients With Co-morbid Schizophrenia and Substance Use Disorders. American Journal of Psychiatry, 158(10), 1706-1713.

Barrowclough, C., Ward, J., Wearden, A. & Gregg, L. (2005). Expressed Emotion and Attributions in Relatives of Schizophrenia Patients With and Without Substance Misuse. Social Psychiatry and Psychiatric Epidemiology, 40(11), 884-91.

Best, D., Campbell, A. & O'Grady, A. (2006). First Annual User Satisfaction Survey. London: National Treatment Agency for Substance Misuse.

[http://www.nta.nhs.uk/publications/documents/nta\\_nta\\_user\\_satisfaction\\_survey\\_2005\\_rb18.pdf](http://www.nta.nhs.uk/publications/documents/nta_nta_user_satisfaction_survey_2005_rb18.pdf)

Best, D., Day, E. & Morgan, B. (2006). Addiction Careers and the Natural History of Change. London: National Treatment Agency for Substance Misuse Research Briefing.

[http://www.nta.nhs.uk/publications/documents/nta\\_addiction\\_careers\\_natural\\_history\\_change\\_2006\\_rb20.pdf](http://www.nta.nhs.uk/publications/documents/nta_addiction_careers_natural_history_change_2006_rb20.pdf)

Brook, J.S., Cohen, P. & Brook, D. (1998). Longitudinal Study of Co-occurring Psychiatric Disorders and Substance Use. Journal of the American Academy of Child and Adolescent Psychiatry. 37, 322-330.

Cantwell, R. (2003). Substance Use and Schizophrenia: Effects on Symptoms, Social Functioning and Service Use. British Journal of Psychiatry, 182, 324-329.

Childline (1997). Beyond the Limit: Children who Live with Parental Alcohol Abuse. London: Childline.

Cleary, M., Hunt, G.E., Matheson, S.L., Siegfried, N. & Walter, G. (2008). Psychosocial Interventions for People with Both Severe Mental Illness and Substance Misuse. Cochrane Database of Systematic Reviews, Issue 1. Art. No.: CD001088. DOI: 10.1002/14651858.CD001088.pub2.

[http://mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD001088/pdf\\_fs.html](http://mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD001088/pdf_fs.html)

Copello, A., Templeton, L. & Powell, J. (2010). The impact of addiction on the family: Estimates of prevalence and costs. Drugs: Education, Prevention, and Policy, 17(1), 63-74.

Day, M. (2004). Keeping the Balance- A Practical Guide to Support Lasting Changes in Drug and Alcohol Abuse. South Warwickshire PCT: Crown Copyright. [www.dh.gov.uk](http://www.dh.gov.uk)

Department of Health (2002). Mental Health Policy Implementation Guide: Dual Diagnosis Good Practice Guide. London: Crown Copyright. [www.dh.gov.uk](http://www.dh.gov.uk)

Department of Health (2006). Models of Care for Alcohol Misusers. London: Crown Copyright.

Dishion, T.J., Veronneau, M.H., Stormshal, E.A. & Kavanagh, K. (2015). Family-centered prevention of adolescent drug abuse: Translational research in a public health framework. Handbook of adolescent drug use prevention: Research, intervention strategies, and practice, 293-867.

Drake, R.E., Mercer-McFadden, C., Mueser, K.T., McHugo, G.J. & Bond, G.R. (1998). Review of Integrated Mental Health and Substance Abuse Treatment for Patients with Dual Disorders. Schizophrenia Bulletin, 24, 589-608.

Drake, R.E., Essock, S.M., Shaner A., Carey, K.B., Minkoff, K., Kola, L., Lynde, D., Osher, F.C., Clark, R.E. & Rickards, L. (2001). Implementing Dual Diagnosis Services for Clients with Severe Mental Illness. Psychiatric Services, 52, 469-476.

DrugScope (2000). Drug Misuse and Mental Health: Learning Lessons on Dual Diagnosis. All Party Parliamentary Drugs Misuse Group. A [www.drugscope.org.uk](http://www.drugscope.org.uk)

Family & Friends (2007). Information for Family and Friends Affected by Others' Substance Misuse. Family and Friends Registered Charity no: 1024360, Warwick, UK. [www.familyfriends-drugs.org.uk](http://www.familyfriends-drugs.org.uk)

Fals-Stewart, W., Birchler, G.R. & O'Farrell, T.J. (1996). Behavioural Couples Therapy for Male Substance Abusing Patients. Effects on Relationships Adjustment and Drug Using Behaviour. Journal of Consulting and Clinical Psychology, 64 (5), 959- 972.

Fals-Stewart, W. & O'Farrell, T.J. (2003). Behavioural Family Counseling and Naltrexalone for Opioid Dependiant Patients. Journal of Consulting and Clinical Psychology, 71 (3), 432-442.

Gournay, K., Sandford, T., Johnson, S. & Thornicroft, G. (1997). Dual Diagnosis of Severe Mental Health Problems and Substance Abuse/Dependence: a Major Priority for Mental Health Nursing. Journal of Psychiatric Mental Health Nursing, 4, 89-95.

Gottlieb, J.D., Mueser, K.T. & Glynn, S.M. (2012). Family therapy for schizophrenia: Co-occurring psychotic and substance use disorders. Journal of Clinical Psychology: In session, 68 (5), 405 – 501.

Haddock, G., Barrowclough, C., Tarrier, N., Moring, J., O'Brien, R., Schofield, N., Quinn, J., Palmer, S., Davies, L., Lowens, I., McGovern, J. & Lewis, S. (2003). Cognitive-Behavioural Therapy and Motivational Intervention for Schizophrenia and Substance Misuse. 18-Month Outcomes of a Randomised Controlled Trial. British Journal of Psychiatry, 183, 418-26.

- Hendriks, V., Van de Schee, E., Blanken, P. (2011). Treatment of adolescents with a cannabis use disorder: Main findings of a randomized controlled trial comparing multidimensional family therapy and cognitive behavioral therapy in The Netherlands. Drug and Alcohol Dependence, 119 (1-2), 64-71.
- Hickie, I.B., Koschera, A., Davenport, T.A., Naismith, S.L. & Scott, E.M. (2001). Comorbidity of Common Mental Disorders and Alcohol or Other Substance Misuse in Australian General Practice. The Medical Journal of Australia, 175, S31-6.
- Hodgins, S., Tengström, A., Bylin, S., Göranson, M., Hagen, L., Janson, M., Larsson, A., Lundgren-Andersson, C., Lundmark, C., Norell, E. & Pedersen, H. (2007). Consulting for Substance Abuse: Mental Disorders Among Adolescents and Their Parents. Nordic Journal of Psychiatry, 61(5), 379-86.
- Holland, M. (1998). Substance Use and Mental Health Problems: Meeting the Challenge. British Journal of Nursing, 7(15), 896-900.
- Horigian, V.E., Espinal, P.S., Alonso, E., Verdeja, R.E., Duan, R., Usaga, I.M., Perez-Lopez, A., Marin-Navarrete, R. & Feaster, D.J. (2016). Readiness and barriers to adopt evidence-based practices for substance abuse treatment in Mexico. Salud Mental, 39(2), 77-84.
- Jacobs, J. (1998). The Links between Substance Misuse and Domestic Violence. London: Alcohol Concern/ Institute for the Study of Drug Dependence (ISSD).
- Johnson, S.D., Stiffman, A., Hadley-Ives, E. & Elze, D. (2001). An Analysis of Stressors and Co-morbid Mental Health Problems that Contribute to Youth's Paths to Substance-Specific Services. Journal of Behavioral Health Services and Research, 28(4), 412-26.
- Johnson, S., Thornicroft, G., Afuwape, S., Leese, M., White, I.R., Hughes, E., Wanigaratne, S., Miles, H. & Craig, T. (2007). Effects of Training Community Staff in Interventions for Substance Misuse in Dual Diagnosis Patients with Psychosis (COMO study): Cluster Randomised Trial. The British Journal of Psychiatry, 191(5), 451 – 452.
- Kamali, M., Kelly, L., Gervin, M., Browne, S., Larkin, C. & O'Callaghan, E. (2000). The Prevalence of Comorbid Substance Misuse and Its Influence on Suicidal Ideation Among In-Patients with Schizophrenia. Acta Psychiatrica Scandinavica, 101(6), 452-6.
- Kamali, M., Kelly, L., Gervin, M., Browne, S., Larkin, C. & O'Callaghan, E. (2001). Psychopharmacology: Insight and Comorbid Substance Misuse and Medication Compliance Among Patients with Schizophrenia. Psychiatric Services, 52(2), 161-3.
- Kavanagh, D. & Barrowclough, C. (2004). Substance misuse in early psychosis: Motivational interventions and individual and family cognitive behaviour therapy. Schizophrenia Research, 70 (1).
- Klostermann, K. (2016). The Oxford handbook of substance use and substance use disorders: Oxford University Press.
- Kumpfer, K.L. & Hansen, W.B. Family-based prevention programs. Parenting and teen drug use: The most recent findings from research, prevention, and treatment, p.166-505.

Lubman, D.I., Hides, L. & Jorm, A.F. (2007). Beliefs of Young People and Their Parents About the Harmfulness of Alcohol, Cannabis and Tobacco for Mental Disorders. The Medical Journal of Australia, 187(5), 266-9.

Mason, P. & Bennett, G. (2004). Promoting Safer Drinking. A Briefing Paper for Drug Workers. London: National Treatment Agency for Substance Misuse. [http://www.nta.nhs.uk/publications/documents/nta\\_promoting\\_safer\\_drinking\\_2004\\_drdaolcol.pdf](http://www.nta.nhs.uk/publications/documents/nta_promoting_safer_drinking_2004_drdaolcol.pdf)

Mason, W.A. & Spoth, R.L. (2012). Sequence of alcohol involvement from early onset to young adult alcohol abuse: Differential predictors and moderation by family-focused preventive intervention. Addiction, 107, 2137-2148.

Meis, L.A., Griffin, J.M., Greer, N., Jensen, A.C., Macdonald, R., Carlyle, M., Rutks, I. & Wilt, T.J. (2013). Couple and family involvement in adult mental health treatment: a systematic review. Clinical Psychology Review, 33 (2), 275 – 286.

Meyer, R. (1995). Clinical Guide to Alcohol Treatment: the Community Reinforcement Approach, New York, Guildford press.

McCrary, D.S. & Hay, W. (1987). Coping with Problem Drinking in the Family. In Orford, J. (Ed) Coping with Disorder in the Family. Croom Helm, London and Sydney

Mueser, K.T., Glynn, S.M., Cather, C., Zarate, R., Fox, L., Feldman, J., Wolfe, R. & Clark, R.E. (2009). Family Intervention for Co-occurring Substance Use and Severe Psychiatric Disorders: Participant Characteristics and Correlates of Initial Engagement and More Extended Exposure in Randomized Controlled Trial. Addictive Behaviors, 34(10), 867-877

Mueser, K.T., Glynn, S.M., Cathers, C., Xie, H., Zarate, R., Fox Smith, L., Clark, R.E., Gottlieb, J.D., Wolfe, R. & Feldman, J. (2013). A randomized controlled trial of family intervention for co-occurring substance use and severe psychiatric disorders. Schizophrenia Bulletin, 39 (3), 658 – 672.

Munroe, A. & Allan, J. (2011). Can family-focused interventions improve problematic substance use in aboriginal communities? A role for social work. Australian Social Work, 64(2), 169-182

Murray, S. & Shenker, D. (2007). Alcohol and the Family. A Positional Paper from Alcohol Concern. London: Alcohol Concern. [http://www.alcoholconcern.org.uk/files/20061220\\_155831\\_Alcohol%20and%20Families%20-%20AC%20Position%20Paper%20FINAL.pdf](http://www.alcoholconcern.org.uk/files/20061220_155831_Alcohol%20and%20Families%20-%20AC%20Position%20Paper%20FINAL.pdf)

National Treatment Agency for Substance Misuse (2006). Guidance for Local Partnerships on User and Carer Involvement. London: NTA. [http://www.nta.nhs.uk/publications/documents/nta\\_guidance\\_local\\_partnerships\\_user\\_carer\\_involvement.pdf](http://www.nta.nhs.uk/publications/documents/nta_guidance_local_partnerships_user_carer_involvement.pdf)

Onken, L.S., Blaine J.D., Genser, S. & Horton, A.M. (1997). Treatment of Drug-Dependent Individuals with Comorbid Mental Disorders. U.S: National Institute on Drug Abuse. Research Monograph 172. <http://www.drugabuse.gov/pdf/monographs/Monograph172/Monograph172.pdf>

Orford, J., Copello, A., Velleman, R. & Templeton, L. (2010). Family members affected by a close relative's addiction: the stress-strain-coping-support model. Drugs: Education, Prevention, and Policy, 17(1), 36-43.

Orr, L.C., Elliott, L. & Barbour, R.S. (2014). Promoting family-focused approaches within adult drug services: The potential of the 'Senses Framework'. International Journal of Drug Policy, 25(5), 888-896.

Pihkala, H., Dimova-Branstrom, N. & Sandlund, m. (2017). Talking about parental substance abuse with children: Eight families' experiences of beardslee's family intervention. Nordic Journal of Psychiatry, 71(5), 395-401.

Rey, J. & Tennant, C. (2002). Cannabis and Mental Health (letter). British Medical Journal 325, 1183-1184.

Robbins, M.S., Feaster, D.J., Horigian, V.E., Rohrbaugh, M., Shoham, V., Bachrach, K., Miller, M., Burlew, K.A., Hodgkins, C., Carrion, I., Vandermark, N., Schindler, E., Werstlein, R. & Szapocznik, J. (2011). Brief strategic family therapy versus treatment as usual: Results of a multisite randomized trial for substance using adolescents, Journal of Consulting and Clinical Psychology, 79 (6), 713-727.

Rossi Menezes, P. & Ratto, L.R. (2004). Prevalence of Substance Misuse Among Individuals with Severe Mental Illness in São Paulo. Social Psychiatry and Psychiatric Epidemiology, 39(3), 212-7.

Rowe, C.L. (2012). Family therapy for drug abuse: review and updates 2003 – 2010. Journal of Marital and Family Therapy, 38(1), 59-81.

Schofield, N., Quinn, J., Haddock, G. & Barrowclough, C. (2001). Schizophrenia and Substance Misuse Problems: a Comparison Between Patients With and Without Significant Carer Contact. Social Psychiatry and Psychiatric Epidemiology, 36(11), 523-8.

Sevy, S., Robinson, D.G., Holloway, S., Alvir, J.M., Woerner, M.G., Bilder, R., Goldman, R., Lieberman, J. & Kane, J. (2001). Correlates of Substance Misuse in Patients With First-Episode Schizophrenia and Schizoaffective Disorder. Acta Psychiatrica Scandinavica, 104(5), 367-74.

Sims, H. (2006). Families in Focus – England. Adfam, Families Drugs and Alcohol, London. <http://www.adfam.org.uk/>

Sparks, S.N., Tisch, R. & Gardner, M. (2013). Family-centered interventions for substance abuse in Hispanic communities. Journal of Ethnicity in Substance Abuse, 12 (1). 68-81.

Stanton, M.D., Steier, F., Cook, L. & Todd, T.C. (1984). Narcotic Detoxification in a Family and Home Context: Final Report 1980-83. Rockville. MD: National Institute on Drug Abuse Treatment Research Branch, US.

Stanton, M.D. & Shadish, W.R. (1997). Outcome, Attrition and Family-Couples Treatment for Drug Abuse: a Meta-Analysis and Review of Controlled Comparative Studies. Psychological Bulletin, 122, 170-191.



Stanton, M.D., Todd, T.C. & Associates. (1982). The Family Therapy of Drug Abuse and Addiction. New York: Guildford Press.

Velleman, R., Orford, J., Templeton, L., Copello, A., Patel, A., Moore, L., Macleod, J. & Godfrey, C. (2011). 12-month follow-up after brief interventions in primary care for family members affected by the substance misuse problem of a close relative. Addiction Research & Theory, 19(4), 362-374.

Wanigaratne, S., Davis, P., Pryce, K. & Brotchie, J. (2005). The Effectiveness of Psychological Therapies on Drug Misusing Clients. NHS National Treatment Agency for Substance misuse Research Briefing 11, June, NTA, London.  
[http://www.nta.nhs.uk/publications/documents/nta\\_effectiveness\\_psycho\\_therapies\\_2005\\_rb11.pdf](http://www.nta.nhs.uk/publications/documents/nta_effectiveness_psycho_therapies_2005_rb11.pdf)

Watts, M. (2007). Management of Severe Mental Illness and Substance Misuse. Nursing Times, 103(14), 28-9.

Weaver, T., Charles, V., Madden, P. & Renton, A. (2002). Co-morbidity of Substance Misuse and Mental Illness Collaborative Study (COSMIC). Research Summaries for Providers and Commissioners. New Series, August 2004. National Treatment Agency for Substance Misuse and the Department of Health, London.  
[http://www.nta.nhs.uk/publications/documents/nta\\_cosmic\\_survey\\_2002\\_rs1.pdf](http://www.nta.nhs.uk/publications/documents/nta_cosmic_survey_2002_rs1.pdf)

Yandoli, D., Eisler, L., Robbins, C., Mulledy, G. & Dare, C. (2002). A Comparative Study of Family Therapy in the Treatment of Opiate Users in a London Drug Clinic. Journal of Family Therapy, 24 (4), 402-422.

Yuen, E. & Toumbourou, J.W. (2011). Does family intervention for adolescent substance use impact parental wellbeing? A longitudinal evaluation. Australian and New Zealand Journal of Family Therapy, 32(3), 249-263.

Zammit, S., Allebeck, P., Andreasson, S., Lundberg, I. & Lewis, G. (2002). Self Reported Cannabis Use as a Risk Factor for Schizophrenia in Swedish Conscripts of 1969: Historical Cohort Study. British Medical Journal, 325, 1199-1201.

## Websites

- **Addaction** a drug and alcohol treatment charity [www.addaction.org.uk](http://www.addaction.org.uk)
- **Adfam** a National organisation working with and for families affected by drugs and alcohol [www.adfam.org.uk](http://www.adfam.org.uk)
- **Alcohol Concern** a national agency on alcohol misuse and also provides information related to mental health. [www.alcoholconcern.org.uk](http://www.alcoholconcern.org.uk)
- **Dual Diagnosis Online Support Group** (UK) a community of patients, family members and friends dedicated to dealing with dual diagnosis together. <http://www.mdjunction.com/dual-diagnosis>
- **Dual Diagnosis Forum** (US) provides an opportunity for networking, idea sharing and reflection among people who provide support to others facing the dual diagnosis. <http://greg.quuxuum.org/dual/>
- **Mental Health America** 'Factsheet: Dual Diagnosis' <http://www.nmha.org/go/information/get-info/co-occurring-disorders/dual-diagnosis>
- **Mind's** leaflet 'Understanding Dual Diagnosis' <http://www.mind.org.uk/NR/rdonlyres/29C42412-155B-4DFF-A950-4AFE2E10FDB3/0/Understandingdualdiagnosis2007.pdf>
- **Rethink's** dual diagnosis leaflet for families, friends and carers <http://www.rethink.org/dualdiagnosis/pdfs/Leaflet.pdf>
- **The Retreat York**, offers a dual diagnosis inpatient service which offers simultaneous treatment for substance misuse and mental health problems. <http://www.theretreatyork.org.uk/ourservices.php?ref=23>
- **Schizophrenia.com** provides in-depth information, support and education related to schizophrenia. This website includes Information on Schizophrenia and cannabis <http://www.schizophrenia.com/prevention/streetdrugs.html>
- The **Substance Abuse and Mental Health Services Administration (SAMHSA)** is an Agency of the US Department of Health and Human Services. It is the federal agency charged with improving the quality and availability of prevention, treatment, and rehabilitative services in order to reduce illness, death, disability, and cost to society resulting from substance abuse and mental illnesses. <http://www.samhsa.gov/index.aspx>
- **Swanswell Charitable Trust** Warwickshire, UK. A leading provider of services which help and support people in tackling their alcohol and/or drug related difficulties, and enable them to lead more independent and fulfilling lives. <http://www.swanswelltrust.org/>